

SEQUESTRATION RATE OF HEAVY METAL CONTAMINANTS USING *RICCIA FLUITANS* AS POTENTIAL PHYTOREMEDIATION AGENT IN POLLUTED AQUATIC ECOSYSTEM

Rashidi Othman¹ --- Nurul Azlen Bt Hanifah^{2*} --- Razanah Ramya³ --- Farah Ayuni Bt Mohd Hatta⁴ --- Wan Syibrah Hanisah Bt Wan Sulaiman⁵ --- Maheran Bt Yaman⁶ --- Zainul Mukrim Bin Baharuddin⁷

^{1,2,3,4,5}International Institute for Halal Research and Training (INHART), Herbarium Unit, ^{6,7}Department of Landscape Architecture, Kulliyah of Architecture and Environment Design, International Islamic University Malaysia, Kuala Lumpur, Malaysia

ABSTRACT

The application of green technology is the combination of the environmental science and technology in which helps the improvement and application of products, apparatus as well as the system to safeguard the natural resources and the environment. Aquatic plants have been increasingly being utilized as remarkably efficient and responsive apparatus for diagnosing and predicting environmental stresses. Thus, this research aimed to assess *Riccia fluitans* capability and potential as a phytoremediation agent to remediate heavy metal contamination in an unhealthy aquatic ecosystem. Subsequently, *Riccia fluitans* was exposed to three types of heavy metals (Mn, Zn and Pb) at three different concentrations (1.0, 2.0 and 5.0 mg L⁻¹) at four different periods of time (week 1 to week 4). All heavy metals were analyzed through HACH DR5000 spectrophotometer. As a result, among five heavy metals tested, *Riccia fluitans* revealed as a good agent to sequester Mn, Zn and Pb respectively. The analysis affirmed the accumulation of different metals within the plant and an equally lessen of metals in the water. An eloquent equivalence between type of heavy metals and concentration, period of time and *Riccia fluitans* were attained. As a conclusion, *Riccia Fluitans* is proven to be a potential biosequester agent for polluted aquatic environment.

Keywords: Green technology, Phytoremediation, Heavy metals, Aquatic ecosystem, Biosequester agent, Polluted environment.

1. INTRODUCTION

An increase in industrial activities is giving birth to a severely injured of environmental pollutions and corroded some aquatic ecosystems correlated with heavy metal accumulation in biota and flora (Thomson *et al.*, 2013). Since the beginning of the industrial revolution, toxic metal pollution of the biosphere has become more exacerbated until causing greater environmental

and health problem. Usability of freshwater runoff had reached nearly 50% and expected to increase another 20% by 2025 (Haris and Wan Maznah, 2008). Beneficial of river has levelled off in the environment due to a poor planning. The monitoring of river water quality is under the responsibility of the Department of Environment Malaysia. Declination shown in the aquatic environment is particularly due to dam construction, river diversions, heavy pollution loads and other habitat changes (Postel *et al.*, 1996).

In fact, Malaysian waters suffered with 30 to 100 times more pathogens, heavy metals and poisons from industry and agriculture above than government standards permit (Hinrichsen and Tacio, 2002).

Macrophytes species are undeniably fitting as an ideal biomonitor organism based on the criteria listed. They are motionless, visible to the naked eye, easy to handle and collectible, easy to find on the site, accumulate and mobilize contaminants and nutrients in their tissues and signifies environmental stresses (Milan *et al.*, 2006). Furthermore, some of aquatic plant species have an outstanding ability in bioconcentration which, hence elevates more nutrients and heavy metal accumulations. In some cases, plant tissues are able to concentrate high amounts in their tissues, resulting in the significant availability of those elements in the surrounding environment (Milan *et al.*, 2006). Thus, macrophytes are applicable as a biosequester agent in aquatic environments. Through a right selection of wetland plant species benefits in greater removal trace element.

Free floating plants are one of the aquatic plant species categories. Like other categories, free floating plants are also capable to accumulate and uptake metal from waste water. One of the family floating plants that assuring results in removing metal from industrial waste matter is Lemnaceae. The term 'Duckweed' associates to this group (Miranda *et al.*, 2000). Duckweed has been diagnosed to remove nutrients and organic matter, suspended solids, various metals (As, Cd, Cr, Cu, Ni, Pb, Fe, Au, and Zn) (Miranda *et al.*, 2000; Kara *et al.*, 2003; Sasmaz and Obek, 2009; Rahman and Hasegawa, 2011; Materazzi *et al.*, 2012). The water hyacinth (*Eichhornia crassipes*), is among the most commonly cited species in purifying polluted waters. The plant is speedy growth and able to accumulate nutrients and metals as well. Furthermore, the plants have tested and potentially remove N, P, As, Cd, Cn, Zn and Pb (Brooks and Robinson, 1998; Ebel *et al.*, 2007; Mishra and Tripathi, 2008; Mishra *et al.*, 2008; Mahamadi and Nharingo, 2010; Rahman and Hasegawa, 2011). The potential use and capacity of *Salvinia natans* for the removal of metals such as Cu, As, Hg, Ni, Fe, Pb, Zn and Cr from industrial effluents was studied and tested out (Sen and Bhattacharyya, 1994; Dhir and Srivastava, 2013; Lima *et al.*, 2014). The plants are also having a high growth rate and metal tolerance capacity to high concentrations (Mandal *et al.*, 2013) The current practices for heavy metal treatment are using reverse-osmosis, ion-exchange,

electrodialysis and adsorption. Majorities are consuming high cost, energy demanding and metal specific (Mishra and Tripathi, 2008) which led to environmental degradation.

Conventional remediation technologies are based on civil and chemical engineering technologies which includes vastly and diversified of physical, thermal and chemical treatments (Hamlin, 2002). Most of metals and radionuclides could not be eliminated from the environment through chemical and biological transformation. The degree of toxicity can be levelled off by through manipulating their speciation, but not in degradation, they are still perpetual in the environment and can accumulate in soil and aquatic environment.

Phytoremediation intends to eliminate heavy metals or other contaminants from soils or waters using plant which granting cost efficiency and environmental sustainability. The plant is used as a hyperaccumulator with exceptional metal-accumulating capacity to extract metals from the soil (Zhao *et al.*, 2003; Gratão *et al.*, 2005). The concept of utilizing plants to immobilize metal and toxic pollutants from polluted environments (soils, waste, water, etc.) is applied nearly about 300 years ago in the treatment of waste water (Gratão *et al.*, 2005) Nonetheless, an early 1980's, there is no scientific study and development of suitable plants were taken in the place. Phytoremediation unmasks plant's unique biological mechanisms for human benefit. Phytoremediation unmasks plant's unique biological mechanisms for human benefit. There are four sections of this technology, which associate to toxic metal remediation of soil and water.

A rising degree and seriousness of environmental problems related to water bodies forces an idea of the present study to find more species that capable of cleaning up aquatic environments contaminated with metals. Hence, the aim of the research is to assess the potential and capability of study species which is *Riccia fluitans* as a potential biosequester agent for polluted aquatic environment. Also, to find out the concentration of Mn, Zn and Pb within the plant in water.

2. MATERIALS AND METHODS

2.1. Plant collection

Riccia fluitans is a liverwort of the genus *Riccia L.* was hand-picked from the canal, located in Taman Tasik Taiping, Perak Darul Ridzuan, Malaysia. The plants were brought back to the laboratory in clean plastic bags for further to study the degree of heavy metal uptake by *Riccia fluitans*. Plants were carefully washed using tap water and then rinse with distilled water, to remove visible debris (Sadler and Rynja, 1992) Then, the process of drying the sample was carefully done using filter papers (Qualitative circles 110mm) for 10 minutes each batch. Dry weights of the plants were registered with a digital balance a (sensitivity of 0.01). Later, they were maintained in a container below an ordinary fluorescent bulb.

2.2. Heavy Metals Preparation

About 1.0 g (fresh weight) of *Riccia fluitans* was floated in 40mL of metal salts (pH 4.5-5.0) in a sterile container (60mL/48mm) in a laboratory conditions. The initial concentrations for individual experiments were varied from 1.0, 2.0 and 5.0 mg/L and controlled of each metal prepared by using $\text{MnSO}_4\text{H}_2\text{O}$, $\text{ZnSO}_4\cdot 7\text{H}_2\text{O}$ and $\text{Pb}(\text{NO}_3)_2$. Each treatment was prepared in ten replicates; a total of 33 containers was involved in this experiment (including control for each concentration). Control treatment (metal without plants) was needed to correlate the results and to confirm the effect of metal ions on *Riccia fluitans* growth. The concentrations of heavy metals in water samples were determined using MERCK standard to provide calibration and quality assurance for each analytical batch. From each tube, samples were measured and collected weekly, and the study was conducted for 4 weeks. All the prepared treatment was maintained at 25°C during the 30 days of Zn, Pb and Mn exposure.

2.3. Heavy Metals Removal

Removal was determined by quantifying the concentration of metal left in the medium after immersion of plants. The plants were taken out from the treatment metal, meanwhile the treatment metal itself was filtered using filter papers (Qualitative circles 110mm). The concentration of heavy metals was determined by DR5000 using reagents by MERCK following each of the elements.

2.4. Statistical Analysis

Analysis of variance (ANOVA) was calculated to test the validity of the data and the significance of the variation in the data of three heavy metals studied Fe, Mn and Cu in different concentration for *Riccia fluitans* at different period of time (week 1 - week 4).

3. RESULTS AND DISCUSSION

Analysis of variance showed significant difference ($p > 0.001$) between heavy metals uptake, concentration ranges (1.0 mg l^{-1} , 2.0 mg l^{-1} and 5.0 mg l^{-1}) and period of time (week 1 until week 4). Based on the results obtained (Figure 1), heavy metals sequestration rate by *Riccia fluitans* were observed on reduction of concentration of Fe, Mn and Cu in water samples. This study documents the capability of *R. fluitans* to remediate was higher by increasing period of time (week 4) at different concentration ranges except for 1 mg l^{-1} , of Mn and Cu and this is in agreement with Mishra *et al.* (2008) and Dhir and Srivastava (2011). In contrast, the emergent aquatic plant species are usually sequestered lower amounts of metals than submerged aquatic species (Kamal *et al.*, 2004). Species such as *Centella asiatica* and *E. crassipes* had a maximum removal of Cu in solution about 99.6% (Mokhtar *et al.*, 2011). In another study, *Lemna minor* showed a resistant to environmental toxicity and rapidly with growing. (Miranda *et al.*, 2000)

On the other hand, the removal efficiencies were increased from 1.0 mg^l⁻¹ up to 5.0 mg^l⁻¹ for all heavy metals studied however certain heavy metals were fluctuated (Figure 1). Both Fe and Mn were decreased at 2 mg^l⁻¹, surprisingly were increased at 5 mg^l⁻¹ except for Cu. As for example in this study showed for Mn at 1 mg^l⁻¹ were 0.69 mg^l⁻¹ at week 1, increasing up to 0.19 mg^l⁻¹ respectively. Cu is an essential micronutrients for plant metabolism, however the excessive can effect on physiological and biochemical processes (Li and Xiong, 2004; Mazen, 2004). At high concentration of Cu, *L. gibba* growth was inhibited due to toxicity (Khellaf and Zerdaoui, 2010). In contrast, *Hydrilla verticillata*, *Elodea Canadensis* and *Salvinia natans* showed toxicity to Cu and Ni also after 5 days (Begum and Hari Krishna, 2010). The experimental plants showed good performance in removing the metal even they were able to remove <50 % of metals solution.


4. CONCLUSION

In conclusion, *Riccia fluitans* was tested for concurrent removal of three heavy metals Fe, Mn and Cu. The aquatic plants proved as effective in remediate and sequester of these heavy metals at three concentration ranges (1.0 mg^l⁻¹, 2.0 mg^l⁻¹ and 5.0 mg^l⁻¹). The high correlation between a period of time and types of heavy metals with different concentration indicates that this plant can be effectively used for removal of heavy metals from solution of different heavy metals.


5. ACKNOWLEDGMENT

The authors would like to thank the Ministry of Higher Education (MOHE) and International Islamic University Malaysia (IIUM) for the Research Grant RACE140-020-0018 and FRGS13-052-0293-.

A.



B.



C.




Figure-1. Assessment of Fe, Mn and Cu sequestration rate by *Riccia fluitans* for 1.0 mg^l-1, 2.0 mg^l-1 and 5.0 mg^l-1 at different incubation period at week 1, week 2, week 3 and week 4

- A. Fe, Mn and Cu sequestration rate for 1.0 mg^l-1 at week 1 until 4
- B. Fe, Mn and Cu sequestration rate for 2.0 mg^l-1 at week 1 until 4
- C. Fe, Mn and Cu sequestration rate for 5.0 mg^l-1 at week 1 until 4

REFERENCES

Begum, A. and S. Hari Krishna, 2010. Bioaccumulation of trace metals by aquatic plants. International Journal of Chem. Tech. Research, 2(1): 250-254.

Brooks, R.R. and B.H. Robinson, 1998. Aquatic phytoremediation by accumulator plants. Plants that hyperaccumulate heavy metals: Their role in phytoremediation, microbiology, archaeology, mineral exploration and phytomining. New York: CAB International. pp: 203-226.

Dhir, B. and S. Srivastava, 2011. Heavy metal removal from a multi-metal solution and wastewater by *Salvinia natans*. Ecological Engineering, 37(6): 893-896.

Dhir, B. and S. Srivastava, 2013. Heavy metal tolerance in metal hyperaccumulator plant, *salvinia natans*. Bulletin of Environmental Contamination and Toxicology, 90(6): 720-724.

- Ebel, M., M.W. Evangelou and A. Schaeffer, 2007. Cyanide phytoremediation by water hyacinths (*Eichhornia Crassipes*). *Chemosphere*, 66(5): 816-823.
- Gratão, P.L., M.N.V. Prasad, P.F. Cardoso, P.J. Lea and R.A. Azevedo, 2005. Phytoremediation: Green technology for the clean up of toxic metals in the environment. *Brazilian Journal of Plant Physiology*, 17(1): 53-64.
- Hamlin, R.L., 2002. Phytoremediation literature review. *Umass. Edu.* 00-00.
- Haris, H. and W.O. Wan Maznah, 2008. The effects of tidal events on water quality in the coastal area of petani river basin, Malaysia. *International Conference on Environmental Research and Technology (ICERT 2008)*; Park Royal Penang, Malaysia. pp: 595-599.
- Hinrichsen, D. and H. Tacio, 2002. The coming freshwater crisis is already here. Finding the source: The linkages between population and water. Washington, DC: Woodrow Wilson International Center for Scholars, Environmental Change and Security Project. pp: 1-26.
- Kamal, M., A.E. Ghaly, N. Mahmoud and R. Cote, 2004. Phytoaccumulation of heavy metals by aquatic plants. *Environment International*, 29(8): 1029-1039.
- Kara, Y.E.Ş.I.M., D. Basaran, I. Kara, A. Zeytinluoglu and H. Genç, 2003. Bioaccumulation of nickel by aquatic macrophyta *lemna minor* (Duckweed). *Int. J. Agr. Biol.*, 5(3): 281-283.
- Khellaf, N. and M. Zerdaoui, 2010. Growth, photosynthesis and respiratory response to copper in *lemna minor*: A potential use of duckweed in biomonitoring. *Iran J. Environ. Health Sci. Eng.*, 7(2): 299-306.
- Li, T.Y. and Z.T. Xiong, 2004. A novel response of wild type of duckweed (*Lemna Paucicostata* Hegelm.) to heavy metals. *Environ. Toxicol.*, 19(2): 95-102.
- Lima, L.K., J.F. Silva, M.G. Da Silva and M.G. Vieira, 2014. Lead biosorption by *salvinia natans* biomass: Equilibrium study. *Chemical Engineering*: 38.
- Mahamadi, C. and T. Nharingo, 2010. Utilization of water hyacinth weed (*Eichhornia Crassipes*) for the removal of Pb (II), Cd (II) and Zn (II) from aquatic environments: An adsorption isotherm study. *Environmental Technology*, 31(11): 1221-1228.
- Mandal, C., N. Ghosh, S. Maiti, K. Das, S. Gupta, N. Dey and M.K. Adak, 2013. Antioxidative responses of *salvinia* (*Salvinia Natans* Linn.) to aluminium stress and it's modulation by polyamine. *Physiology and Molecular Biology of Plants*, 19(1): 91-103.
- Materazzi, S., S. Canepari and S. Aquili, 2012. Monitoring heavy metal pollution by aquatic plants. *Environmental Science and Pollution Research*, 19(8): 3292-3298.
- Mazen, A.M.A., 2004. Accumulation of four metals in tissues of *corchorus olitorius* and possible mechanisms of their tolerance. *Biol. Plant*, 48(2): 267-272.
- Milan, B., P. Slobodanka, S. Živko and K. Borivoj, 2006. Macrophytes as phytoindicators and potential phytoremediators in aquatic ecosystems. In th International Conference: International Austrian Committee Danube Research, Vienna.
- Miranda, G., A. Quiroz and M. Salazar, 2000. Cadmium and lead removal from water by the duckweed. *Lemna gibba* (Lemnaceae). *Hidrobiologica (Iztapalapa)*. Iztapalapa, 10(1): 7-12.
- Mishra, V.K. and B.D. Tripathi, 2008. Concurrent removal and accumulation of heavy metals by the three aquatic macrophytes. *Bioresource Technology*, 99(15): 7091-7097.

- Mishra, V.K., A.R. Upadhyaya, S.K. Pandey and B.D. Tripathi, 2008. Heavy metal pollution induced due to coal mining effluent on surrounding aquatic ecosystem and its management through naturally occurring aquatic macrophytes. *Bioresource Technology*, 99(5): 930-936.
- Mokhtar, H., N. Morad and F.F. Ahmad Fizri, 2011. Hyperaccumulation of copper by two species of aquatic plants. *International Conference on Environmental Science and Engineering*, 8: 115-118.
- Postel, S.L., G.C. Daily and P.R. Ehrlich, 1996. Human appropriation of renewable fresh water. *Science*, 271(5250): 785-787.
- Rahman, M.A. and H. Hasegawa, 2011. Aquatic arsenic: Phytoremediation using floating macrophytes. *Chemosphere*, 83(5): 633-646.
- Sadler, R. and G. Rynja, 1992. Preservation, storage, transport, analysis and reporting of water samples, Queensland government chemical laboratory report, Series No. 12. Brisbane: Queensland Government Publishers. U.N.
- Sasmaz, A. and E. Obek, 2009. The accumulation of arsenic, uranium, and boron in *Lemna gibba* L. exposed to secondary effluents. *Ecological Engineering*, 35(10): 1564-1567.
- Sen, A.K. and M. Bhattacharyya, 1994. Studies of uptake and toxic effects of Ni (II) on *salvinia natans*. *Water, Air, and Soil Pollution*, 78(1-2): 141-152.
- Thomson, I.S.I., K.S. Uka, Chukwuka and C. Afoke, 2013. Heavy metal accumulation by *Telfairia occidentalis* hook f. Grown on waste dumpsites in South-Eastern Nigeria. *Research Journal of Environmental Toxicology*, 7(1): 47-53.
- Zhao, F.J., J.R. Wang, J.H.A. Barker, H. Schat, P.M. Bleeker and S.P. McGrath, 2003. The role of phytochelatins in arsenic tolerance in the hyperaccumulator *pteris vittata*. *New Phytologist*, 159(2): 403-410.

BIBLIOGRAPHY

- Dalwani, D., S. Dixit and S. Dote, 2008. Evaluation of uptake rate of heavy metals by *eichhornia crassipes* and *Hydrilla verticillata*. *Environmental Monitoring Assessment*, 169 (1-4): 367-374.
- Dhir, B. and S. Srivastava, 2012. Disposal of metal treated *Salvinia* biomass in soil and its effect on growth and photosynthetic efficiency of wheat. *International Journal of Phytoremediation*, 14(1): 24-34.
- LI, Q., X.Y. GUO, X.H. XU, Y.B. ZUO, D.P. WEI and Y.B. MA, 2012. Phytoavailability of copper, zinc and cadmium in sewage sludge-amended calcareous soils. *Pedosphere*, 22(2): 254-262.
- Mishra, V.K. and B.D. Tripathi, 2009. Accumulation of chromium and zinc aqueous solutions using water hyacinth (*Eichhornia Crassipes*). *Journal of Hazardous Material*, 164(2-3): 1059-1063.

Views and opinions expressed in this article are the views and opinions of the author(s), International Journal of Sustainable Energy And Environmental Research shall not be responsible or answerable for any loss, damage or liability etc. caused in relation to/arising out of the use of the content.